



IL-2 and IL-10 serum levels in HIV-1-infected patients with or without active antiretroviral therapy

Orsilles, Miguel Ángel, Pieri, Elsa Cristina, Cooke, Paula María  and Caula, Cinthya  (2006) *IL-2 and IL-10 serum levels in HIV-1-infected patients with or without active antiretroviral therapy*. *APMIS*, 114 (1). pp. 55-60. ISSN 1600-0463

El texto completo no está disponible en este repositorio.

RESUMEN

We analyzed IL-2 and IL-10 serum levels in 26 HIV-1-infected patients naive of antiretroviral treatment and in 34 patients receiving highly active antiretroviral therapy (HAART). All patients without treatment were asymptomatic. When they were stratified according to levels of CD4+ T cells, IL-2 levels were significantly increased in patients with ≥ 200 CD4+/ μ l and IL-10 levels were significantly increased in patients with...

TIPO DE DOCUMENTO:

Artículo

DOI:

https://doi.org/10.1111/j.1600-0463.2006.apm_108.x

PALABRAS CLAVE:

HAART. HIV. IL-10. IL-2.

TEMAS:

[Q Ciencia](#) > [QD Química](#)

UNIDAD ACADÉMICA:

[Universidad Católica de Córdoba](#) > [Facultad de Ciencias Químicas](#)